


# Age Related Residential Care Vitamin D Assessment Tool



\* Vitamin D supplement brand Vit.D3 contains soya oil; some people are allergic to soya oil and some people with peanut allergy also react to soya oil.

\*\* Vitamin D supplementation should be considered for all patients who receive long-term treatment with any antiepileptic medicine, including the newer agents.<sup>1</sup> Falls risk is increased, and vitamin D deficiency common, in patients with epilepsy.<sup>2</sup>

Developed by Brendan Duck, Clinical Pharmacist, in association with Summerset in the Vines. Approved by: Dr Tim Frendin March 2013  
Updated by Di Vicary, Clinical Pharmacist, reviewed by Brendan Duck and Caroline Chandler, endorsed by Dr Ian Hosford, HBDHB SMO - Older Persons. April 2016. References

1. Medicines and Healthcare products Regulatory Agency. Antiepileptics: adverse effects on bone. 1 April 2009
2. Petty SJ, Wilding H, Wark JD. Osteoporosis Associated with Epilepsy and the Use of Anti-Epileptics – a Review. *Curr Osteoporosis Rep*; 2016(14):54-65
3. bpacnz Prescribing issues associated with anticonvulsant medications for epilepsy. *Best Practice Journal*. November 2009. Available from: <http://www.bpac.org.nz/BPJ/2009/November/anticonvulsants.aspx> (Accessed 8 April 2016)
4. HQSC. Vitamin D and falls: what you need to know, and, further reading. <http://www.hqsc.govt.nz/our-programmes/reducing-harm-from-falls/10-topics/topic-7/>
5. ACC. Vitamin D Prescribing Advice. October 2012
6. Australian Commission on Safety and Quality in Health Care. Falls facts for doctors. Preventing Falls and Harm From Falls in Older People: Best Practice Guidelines for Australian Hospitals 2009